

tumor. Preliminary results suggest that this approach seems a reasonable option for Stage I unilateral Wilms' tumor [5,6].

Francesco Cozzi, MD  
Amalia Schiavetti, MD  
Pediatric Surgery Unit and Pediatric Oncology Unit  
University of Rome "La Sapienza"  
Policlinico Umberto I  
00161, Rome, Italy

## REFERENCES

1. Ritchey ML, Green DM, Thomas PRM, Smith GR, Haase G, Shochat S, Moksness J, Breslow NE: Renal failure in Wilms' tumor patients: a report from the National Wilms' Tumor Study Group. *Med Ped Oncol* 26:75–80, 1996.
2. Morgan WR, Zincke H: Progression and survival after renal-conserving surgery for renal cell carcinoma: experience in 104 patients and extended follow-up. *J Urol* 144:852–885, 1990.
3. Stephens R, Graham SD Jr: Enucleation of tumor versus partial nephrectomy as conservative treatment of renal cell carcinoma. *Cancer* 65:2663–2667, 1990.
4. Herr HW: Partial nephrectomy for renal cell carcinoma with normal opposite kidney. *Cancer* 73:160–162, 1994.
5. Cozzi F, Schiavetti A, Clerico A, Matrunola M, Cozzi DA, Castello MA: Tumor enucleation in unilateral Wilms' Tumor: a pilot study [abstr]. *Med Ped Oncol* 25:313, 1995.
6. Cozzi I, Schiavetti A, Bonanni M, Matrunola M, Cozzi DA, Castello MA: Enucleative surgery for stage I nephroblastoma with a normal contralateral kidney. *J Urol* 156: in press, 1996.

## Letter to the Editor: Response to Letter by Cozzi and Schiavetti

We (the authors) recognize that a number of centers are employing protocols for the use of nephron sparing surgery in children with unilateral nephroblastoma. The purpose of our review was to establish the known risk of renal failure following treatment of Wilms' tumor. The National Wilms' Tumor Study Group has a long-standing commitment to track the late effects of children treated for this disease. The incidence of renal failure that we reported is the best data that is known, to date, in such a large series of patients. It will be many years before all of these patients reach late adulthood and we will be able to determine the exact risk of late clinical renal insufficiency. However, with follow-up ranging up to 25 years, we have noted a very low risk for patients with unilateral tumors.

The letter by Cozzi and Schiavetti referred to the experience with renal cell carcinoma. There is a difference between the latter patients and those with Wilms' tumor. Patients with renal cell carcinoma that are amenable to partial nephrectomy have small tumors at diagnosis. Most Wilms' tumor patients require preoperative chemotherapy to facilitate partial resection. The experience in bilateral tumors suggests that this can be feasible [1]. The National Wilms' Tumor Study Group believes that this is an acceptable treatment for bilateral tumors, given the known increased incidence of renal failure in this group of patients. However, in order to employ this approach for unilateral tumors, one would then have to weigh the small known risk of renal failure in this group vs. the problems with

inaccurate staging following chemotherapy [2,3]. Therefore, we have not considered it necessary to adopt such an approach to date.

Michael L. Ritchey, MD  
Daniel M. Green, MD  
Patrick R.M. Thomas, MD  
Glenda R. Smith, MD  
Gerald Haase, MD  
Stephen Shochat, MD  
Jami Moksness, BS  
Norman E. Breslow, PhD  
Department of Surgery  
The University of Texas  
Houston Medical School  
Houston, TX 77030

## REFERENCES

1. Horwitz, et al: Renal salvage procedures in patients with synchronous bilateral Wilms' tumor: a report from the National Wilms' Tumor Study group. *Journal of Pediatric Surgery* (in press).
2. Tournade MF, Com-Nouge C, Voute PA, Lemerle J, de Kraker J, Delemarre JFM, et al: Results of the sixth International Society of Pediatric Oncology Wilms' tumor trial and study: a risk-adapted therapeutic approach in Wilms' tumor. *J Clin Oncol* 11:1014–1023, 1993.
3. Green DM, Breslow NE, D'Angio GJ: The treatment of children with unilateral Wilms tumor. *J Clin Oncol* 11:1009–1010, 1993.